





Fix the Cat on the operation table.

Take a 4 pieces of 1 m long bandages and each fold in the middle.

Make a loop so the two free ends are similar lenght.

Put each loop on the carpal and metatarsal joints and tye a shoetie knot to the oparation table angles.



Operarion site desinfection – surgical scrub.

Take a box of Spiritus and box of chlorheksidine wipes.

- 1. Scrub from proposed incision line circular to perifery with alcohol gauze swab at least 3 times. Skin should be clean. Never return to the proposed incision line with the same gauze swab. Then replace gauze swab with clean one.
- 2. Then take chlorheksidine wipes and repeat scrubbing as with alcohol gauze swab.
- 3. Last step is to take spiritus gauze swab and repeat action as in the first step.

Remember that total contact time for desinbectants should be at least 5 minutes.



Prepare the necessary equipment and materials in the surgery room

Equipment

- Clipper
- Pulse oximeter
- Operating mattress
- Absorbent sheet

Materials

- Chlorhexidine tampons
- Alcohol swabs
- Gauze ties Wound patch
- Wound protection spray (Aluminum spray)

Sterile surgical materials

- Surgical thread
- Scalpel blade
- Surgical instrument set
- Sterile surgical swabs
- Sterile surgical drape
- Sterile instrument cloth
- Sterile surgical gown

Information before starting the scenario



Information:

FOR THE SUERVISOR:

- 1. Read the information for the supervisor and the leader out loud.
- 2. Observe how the leader performs actions according to the algorithm steps (on the left side).
- 3. Do not comment.
- 4. If the leader performs (tells) the written action mark it.
- 5. The text in gold should not be read aloud (information for supervisor only).
- 6. At the end of the situation, discuss the actions of the leader: what was performed well and what was not performed.

FOR THE LEADER

- 1. The patient should be examined as in a real life situation.
- During the scenario look at the algorithm (●), read the actions out loud (◄), perform the action (◄).

FOR THE ASSISTANT

- 1. You are an assistant during this scenario you are acting as a patient / nurse / relative.
- 2. You have to read aloud the text on the right (text in green) for the leader.
- 3. Read the answer only after the leader asks you.

A clinically healthy sphynx cat has been brought to the clinic for OHE surgery. You take the cat in the portable cage and place it on the table to prepare for the anesthetic injection and the OHE operation. The owner has informed that the cat is aggressive.

In preparation, the veterinarian performs a subcutaneous injection and OHE surgery.


Nr.	Steps of the algorithm (performed by the leader and assistant)	>	Course of the scenario (the supervisor reads out loud - only the text that is marked black)
1.	Feline OHE pre and postoperative care		
2.	Pre-operations care		
3.	Inspect equipment required for surgery		OPEN Prepare the necessary equipment and materials in the surgery room
4.	Take the cat which is in the animal in the portable cage		
5.	Fix the cat for subcutaneous injection		

6.	Cat fixation for s.c. injection		
7.	Ask the owner or a cat is aggressive?		the cat is aggressive
8.	Take the animal fixation cages		
9.	Place the cat in a fixation cage		
10.	An injection is made by a veterinarian		the injection is given by a veterinarian
11.	Wait 10 minutes until the cat is fell asleep		
12.	Check the tone of the hind leg muscles by extending the hind leg		The action is accepted when the cat's hind leg is extended
13.	Did the cat show resistance?		the cat resists
14.	Wait 10 minutes until the cat is asleep		
15.	Check the tone of the hind leg muscles by extending the hind leg		The action is accepted when the cat's hind leg is extended
16.	Did the cat resist?		the cat shows no resistance
17.	Take the cat out of the cage and put on surgery table		
18.	Position the cat ventrodorsal position		action is accepted if the cat is in a dorsoventral position OPEN cat in dorsoventral position
19.	Put on the cat's tongue pulse oximeter		the action is accepted if the cat has a pulse oximeter on its tongue

20.	Attach all 4 legs on the surgical table		the operation is accepted if 4 gauze ties of at least 1 m length are taken and folded in half the operation is accepted if a loop is created at the fold for all 4 gauze tie the action is assumed when the loops are placed over the carpal and metatarsal joints action is accepted if all 4 ties are tied to the table with a simple shoelace knot OPEN Fix the Cat on the operation table.pdf
21.	Prepare operations square		first clean with alcohol-based skin disinfectant surgical swabs second, the surgical site is cleaned with surgical swabs containing chlorhexidine third cleaned with alcohol surgical swabs all cleaning steps are performed circularly from the center to the periphery
22.	Post-operations care		
23.	Clean from blood postoperative wound with hydrogen peroxide and surgical tampons		accepted if the wound is clean of blood
24.	Treat a wound with a wound protective spray		the action is accepted if a wound spray is blown on the wound
25.	Put on the wound patches		the action is accepted if a wound patch is stuck on the postoperative wound
26.	Free the limbs from the fixation links		the action is accepted if all ties from the table to the animal are untied

1			
27.	Remove the pulse oximeter		action taken when the pulse oximeter is removed from the tongue
28.	Put on postoperative collars		accepted if the cat has a postoperative collar
29.	Remove the animal from surgical table		accepted if the cat is taken from the table and put in the cage
30.	Put the cat down in a portable cage, in lateral sleep, facing the cage for the door		accepted if the cat's head is placed against the door of the cage
31.	Check the eyelid reflex		accepted if the eyelid reflex is tested
32.	When the eyelid reflex is positive cats can be left alone		eyelid reflex is positive
33.		End	





© 2022 All rights reserved by "Hybrid Lab". You may not distribute, copy, reproduce or store any portion of the product in the search engines without the copyright owner's express written consent. Version v.1.1. / ALG-HYBVET-000127



Information before starting the scenario



Information:

FOR THE SUPERVISOR:

- 1. Read the information for the supervisor and the leader out loud.
- 2. Observe how the leader performs actions according to the algorithm steps (on the left side).
- 3. Do not comment.
- 4. If the leader performs (tells) the written action mark it.
- 5. The text in gold should not be read aloud (information for supervisor only).
- 6. At the end of the situation, discuss the actions of the leader: what was performed well and what was not performed.

FOR THE LEADER

- 1. The patient should be examined as in a real life situation.
- 2. During the scenario look at the algorithm (), read the actions out loud (), perform the action ().

FOR THE ASSISTANT

- 1. You are an assistant during this scenario you are acting as a patient / nurse / relative.
- 2. You have to read aloud the text on the right (text in green) for the leader.
- 3. Read the answer only after the leader asks you.

A sphynx cat has been brought to the veterinary clinic for OHE surgery. The owner of the animal says that the cat is aggressive.

Nr.	Steps of the algorithm (performed by the leader and assistant)	>	Course of the scenario (the supervisor reads out loud - only the text that is marked black)
1.	Feline OHE pre and postoperative care		
2.	Pre-operations care		
3.	Take a cat with an animal portable cages		
4.	Fix the cat for subcutaneous injection		
5.	Cat fixation for s.c. injection		
6.	Take the cat which is in the animal in the portable cage		
7.	Ask the owner or a cat is aggressive?		the cat is aggressive

8.	Take the animal fixation cages		
9.	Insert the cat into the animal in the fixation cage		
10.	The veterinarian performs injection for the cat		
11.	Wait 10 minutes until the cat is fell asleep		
12.	Check the hind legs muscle tone stretching the leg		
13.	Did the cat show resistance?		The cat shows no resistance
14.	Take the cat out of the cage and put on surgery table		
15.	Position the cat ventrodorsal position		OPEN cat in dorsoventral position
16.	Put on the cat's tongue pulse oximeter		OPEN Pulse oximeter on a cat's tongue
17.	Attach all 4 legs on the surgical table		OPEN Fix the Cat on the operation table.pdf
18.	Prepare operations square		OPEN Departion site desinfection.pdf
19.	Postoperative care		
20.	Post-operations care		
21.	Clean from blood postoperative wound with hydrogen peroxide and surgical tampons		
22.	Treat a wound with a wound protective spray		
23.	Put on the wound patches		

24.	Free the limbs from the fixation links		
25.	Remove the pulse oximeter		
26.	Put on postoperative collars		
27.	Remove the animal from surgical table		
28.	Put the cat down in a portable cage, in lateral sleep, facing the cage for the door		
29.	Check the eyelid reflex		The eyelid reflex is negative
30.	When the eyelid reflex is positive cats can be left alone		
31.		End	